arXiv:1110.2679v1 [cond-mat.str-el] 12 Oct 2011

Hubbard-I approach to the Mott transition

Przemysław R. Grzybowski[1] and Ravindra W. Chhajlany[1]

[1]*Faculty of Physics, Adam Mickiewicz University, Umultowska 85, 61-614 Poznań, Poland*

(Dated: October 13, 2011)

A Hubbard-I like approach is reconsidered in the context of the metal-insulator transition in strongly correlated electron systems. We show that this approach yields a simple treatment of the transition (at half-filling) and the metallic phase (for arbitrary fillings) in a constrained Hubbard model, that is equivalent to a particular case of a Hubbard bond charge Hamiltonian on bipartite lattices. These calculations, which are to our knowledge the first Hubbard-I calculations for this model, yield results which compare well with known results for this model. For half filling, we also compare these results with those obtained using the Gutzwiller approximation. Finally, we propose a candidate variational wave function connecting the Gutzwiller wave function approach with the scheme presented here.

PACS numbers: 71.30+h,71.10.Fd,71.10.Ay

Introduction The metal to Mott insulator (MI) transition, envisaged by Mott [1], is one of the striking effects induced by strong electronic correlations in many electron systems. The Coulomb interaction between electrons U leads to highly correlated ground states in these systems, rendering their description quite challenging. As a result there have been various theoretical attempts at providing a satisfying description of the Mott transition.

The first of these was due to Hubbard [2], who provided a seminal nonperturbative approach – the so-called Hubbard-I (H-I) approximation – to a simplified interacting electron problem described by the Hubbard model. H-I describes both (i) the atomic limit (*i.e.* limit of vanishing bandwidth $W \to 0$) exactly, in particular yielding two atomic levels corresponding to single and double local occupancy, and (ii) the non-interacting case ($U = 0$) exactly, and so held some promise of describing the intermediate physics in a consistent interpolating fashion. For finite bandwidth, the atomic levels broaden into two "dynamic" (sub)bands which are occupation-number and interaction dependent. These are always split by a gap for all $U > 0$ indicating an insulator phase. Unfortunately, a metal-insulator transition is predicted exactly at $U = 0$, associated with gap closure. The situation was somewhat improved in the Hubbard-III approximation [3], where scattering and resonance broadening corrections shift the transition point to $U_c \sim W$. However, this approach does not predict the expected Fermi Liquid properties on the metallic side, [4]. These drawbacks not withstanding, the Hubbard approaches are very important for the conceptual introduction of the Hubbard subbands and also because they give a basic description of the insulating phase.

A complementary approach, starting from the weak coupling limit, is based on the Gutzwiller variational wave function [5], which describes an increasingly correlated Fermi liquid with increasing interaction U. Brinkman and Rice [6] showed, using the so-called Gutzwiller approximation [5], that increase in U is associated with the diminishing of the quasiparticle residue of the Fermi liquid, or equivalently with the increase in quasiparticle effective mass. In this framework, a metal-insulator transition occurs when the effective mass diverges and is thus driven by quasiparticle localisation. The critical value U_c obtained in the (Gutzwiller)-Brinkman-Rice (GBR) approach is much larger than in the Hubbard approach. This essentially weak coupling method gives a good low energy description of the metal, but does not describe the precursors of the Hubbard bands which should plausibly appear on the metallic side.

More involved contemporary methods, such as dynamical mean field theory (DMFT) [7] or Green's function ladder diagram techniques[8], provide a connection between a strongly correlated fermi liquid picture on the one hand, and formation of Hubbard bands on the other. While providing greater insight into the physics at play, these methods do not provide a simple picture characteristic of more analytical techniques. In this paper, we reconsider the Hubbard-I approach as a systematic means of obtaining a picture, complementary to the one obtained using the Gutzwiller approximation, of the Mott transition from the point of view of strong coupling theory. We shall show that there are strongly correlated Hamiltonians for which the Hubbard-I approach yields good qualitative and quantitative results.

Inspection of the Hubbard-I approach Contrary to early expectations, the Hubbard approach, based on a Green's function decoupling scheme, is now seen as a large–U approximation, despite its non perturbative nature. One may therefore suspect that some problems encountered in the Hubbard approach may originate in limitations of strong-coupling perturbation theory.

Consider therefore the Hubbard model: $H = \hat{T} + U\hat{D}$, where $\hat{D} = \sum_i n_{i\uparrow} n_{i\downarrow}$ is the number of doubly occupied sites and $\hat{T} = \sum_{i,j,\sigma} t_{i,j} c^{\dagger}_{i\sigma} c_{j\sigma}$ is the kinetic energy. The large-U perturbational expansion starts from splitting the kinetic energy into three parts

$\hat{T} = \hat{T}_0 + \hat{T}_{+1} + \hat{T}_{-1}$. $\hat{T}_0 = \hat{T}_{UHB} + \hat{T}_{LHB}$ is the double occupancy conserving hopping (commuting perturbation) in the Upper and Lower Hubbard Bands $\hat{T}_{UHB} = \sum_{i,j,\sigma} t_{ij} n_{i\bar{\sigma}} c^{\dagger}_{i\sigma} c_{j\sigma} n_{j\bar{\sigma}}, \hat{T}_{LHB} = \sum_{i,j,\sigma} t_{ij} (1 - n_{i\bar{\sigma}}) c^{\dagger}_{i\sigma} c_{j\sigma} (1 - n_{j\bar{\sigma}})$, corresponding to hopping of projected electrons on doubly and singly occupied sites respectively, and interband hopping terms $\hat{T}_{+1}$ and $\hat{T}_{-1} = \hat{T}^{\dagger}_{+1}$ (off-diagonal perturbation). The perturbational expansion for the Hubbard model is well known and can be performed e.g. using the method of canonical transformations [9, 10] which to second order yields the effective Hamiltonian:

$$H^{(2)}_{\text{eff}} = \hat{T}_0 + U\hat{D} + \frac{1}{U}[\hat{T}_{+1}, \hat{T}_{-1}]. \tag{1}$$

Recall that the perturbation expansion to any order leads to an effective Hamiltonian eliminating mixing between the degenerate subspaces of the unperturbed Hamiltonian $\hat{U}$. Due to this property, the average of operators jointly changing the total number of doubly occupied sites, such as *e.g.* $\langle c_{j\sigma} n_{j\bar{\sigma}} c^{\dagger}_{l\sigma} h_{l\bar{\sigma}} \rangle$,($h_{j\bar{\sigma}} = 1 - n_{j\bar{\sigma}}$ is the on-site hole occupation number), are identically zero leading to the vanishing of the associated Green's functions. Thus in this framework, the single particle Green's function $G(j,l) = \langle\langle c_{j\sigma} | c^{\dagger}_{l\sigma} \rangle\rangle$ decomposes into a sum of two Green's functions $\Gamma^{\sigma}_{lj} = \langle\langle c_{j\sigma} n_{j\bar{\sigma}} | c^{\dagger}_{l\sigma} n_{l\bar{\sigma}} \rangle\rangle$ and $\widetilde{\Gamma}^{\sigma}_{jl} = \langle\langle c_{j\sigma} h_{j\bar{\sigma}} | c^{\dagger}_{l\sigma} h_{l\bar{\sigma}} \rangle\rangle$ related to the propagation of fermionic quasiparticles, in the upper and lower Hubbard bands.

Consider the Hubbard-I approach to the simplest nontrivial level of strong-coupling perturbational expansion, described by the first two terms in the expansion Eq.(1)

$$H_c = \hat{T}_0 + U\hat{D}. \tag{2}$$

In the frequency domain, the equations of motion for the basic Green's function are $\omega\Gamma^{\sigma}_{lj}(\omega) = \frac{1}{2\pi}\langle n_{l\bar{\sigma}} \rangle \delta_{lj} + \langle\langle [c_{l\sigma} n_{l\bar{\sigma}}, H_{\text{c}}] | c^{\dagger}_{j\sigma} n_{j\bar{\sigma}} \rangle\rangle_{\omega}$ and $\omega\widetilde{\Gamma}^{\sigma}_{lj}(\omega) = \frac{1}{2\pi}\langle h_{l\bar{\sigma}} \rangle \delta_{lj} + \langle\langle [c_{l\sigma} h_{l\bar{\sigma}}, H_{\text{c}}] | c^{\dagger}_{j\sigma} h_{j\bar{\sigma}} \rangle\rangle_{\omega}$. We perform the following type of decoupling on the higher order Green's function in the equations for $\Gamma^{\sigma}_{lj}(\omega)$: $\langle\langle c_{l+\delta\sigma} n_{l+\delta\bar{\sigma}} n_{l\bar{\sigma}} | c^{\dagger}_{j\sigma} \rangle\rangle_{\omega} \approx \langle n_{l\bar{\sigma}} \rangle \Gamma^{\sigma}_{l+\delta,j}(\omega)$, $\langle\langle c_{l\sigma} c^{\dagger}_{l\bar{\sigma}} c_{l+\delta\bar{\sigma}} n_{l+\delta\sigma} | c^{\dagger}_{j\sigma} \rangle\rangle_{\omega} \approx \langle c_{l\sigma} c^{\dagger}_{l\bar{\sigma}} \rangle \langle\langle c_{l+\delta\bar{\sigma}} n_{l+\delta\sigma} | c^{\dagger}_{j\sigma} \rangle\rangle_{\omega}$, $\langle\langle h_{l+\delta\sigma} c^{\dagger}_{l+\delta\bar{\sigma}} c_{l\bar{\sigma}} c_{l\sigma} | c^{\dagger}_{j\sigma} \rangle\rangle_{\omega} \approx \langle c_{l\bar{\sigma}} c_{l\sigma} \rangle \langle\langle c^{\dagger}_{l+\delta\bar{\sigma}} h_{l+\delta\sigma} | c^{\dagger}_{j\sigma} \rangle\rangle_{\omega}$, with analogous complementary treatment of the set $\widetilde{\Gamma}^{\sigma}_{lj}(\omega)$. The last two decouplings lead to magnetic and superconducting order parameters are set here to zero, as we are interested in the description of a pure Mott transition. Solving the equations of motion, one obtains the following momentum (or Fourier transformed) Green's functions for the system:

$$\Gamma^{\sigma}_{\boldsymbol{k}}(\omega) = \frac{1}{2\pi} \frac{n_{\bar{\sigma}}}{\omega - (n_{\bar{\sigma}}\epsilon_{\boldsymbol{k}} + U - \mu)} \tag{3}$$

$$\widetilde{\Gamma}^{\sigma}_{\boldsymbol{k}}(\omega) = \frac{1}{2\pi} \frac{1 - n_{\bar{\sigma}}}{\omega - ((1 - n_{\bar{\sigma}})\epsilon_{\boldsymbol{k}} - \mu)} \tag{4}$$

The electron Green's function thus has the characteristic Hubbard two pole form describing two subbands, however the poles here have an elementary form in comparison with the solution for the full Hubbard model [2]. The main result stemming from these Green's functions is that a Mott transition is elevated to finite interaction U. Indeed, for a paramagnetic configuration at half filling $\langle n_{\uparrow} \rangle = \langle n_{\uparrow} \rangle = 1/2$ the lowest energy in the UHB is $U - W/4$ while the highest energy in the LHB is $W/4$ (where W is the free electron bandwidth), hence a gap opens at a critical value $U_c = W/2$. A fundamental drawback of the Hubbard-I approach to the full Hubbard model is that the resultant Green's functions do not conserve the electron-hole symmetry at half-filling. Importantly, the presented solutions Eqs.(3,4) are explicitly electron hole symmetric and do not suffer from this drawback. Accordingly, we show below that the gap opening at $U = W/2$ is associated with a Mott transition only exactly at half-filling. These results provide explicit evidence that the source of problems in the Hubbard-I approach stem from the effects of the $T_{\pm 1}$ terms.

The Mott transition The constrained Hamiltonian H_c Eq.(2) is interesting in itself because it can be viewed as a particular model of extremely correlated electrons. Recall that the term extremely correlated electron systems has recently been introduced by Shastry [11], emphasizing the appearance of noncanonical fermions resulting from the prohibition of double occupancies in the $U = \infty$ limit, which is a special case of symmetry of conserved double occupancies. The model H_c considered here, that allows for double occupancies which are conserved and thus describes hopping of noncanonical (constrained) fermions $c_{j\sigma} n_{j\bar{\sigma}}$ and $c_{j\sigma} h_{j\bar{\sigma}}$ can therefore be considered a generalization of the problem of extremely correlated fermions for general interaction strength U. Note that the model H_c contains the $U \to \infty$ limit of the Hubbard model. Even for small U and low fillings, we discuss below that the ground states of H_c correspond to the ground states of the $U = \infty$ Hubbard model.

We now consider the details of the Hubbard-I description of the zero temperature Mott transition in the model H_c of extremely correlated electrons. The analysis is carried out only for the paramagnetic phases, with $n_{\uparrow} = n_{\downarrow}$.

From the band structure of the one particle Green's function, it is evident that the insulator phase exists only if $U > U_c$ and the lower band is completely filled while the upper band is empty. The Green's functions Eqs.(3,4) lead to the following equation for the number of electrons of a given spin species:

$$n_{\sigma} = (1 - n_{\bar{\sigma}}) \int_{-W}^{W} f((1 - n_{\bar{\sigma}})\epsilon - \mu)\rho(\epsilon)\,\mathrm{d}\epsilon$$

$$+ n_{\bar{\sigma}} \int_{-W}^{W} f(n_{\bar{\sigma}}\epsilon + U - \mu)\rho(\epsilon)\,\mathrm{d}\epsilon \tag{5}$$

where $f(\dots)$ is the Fermi-Dirac distribution and $\rho(\epsilon)$ is the density of states. The boundaries of the insulator phase are obtained, independently of the lattice, when either $\mu = (1 - n_{\bar{\sigma}})W/2$ is at the end of the lower band or $\mu = -n_{\bar{\sigma}}W/2 + U$ is at the beginning of the upper band. For both these cases, at zero temperature, Eq.(5) reduces to $n_\sigma = 1 - n_{\bar{\sigma}}$, showing that the transition only occurs at half-filling. The jump in μ at half-filling $n_\uparrow = n_\downarrow = 1/2$ reduces to zero at $U = W/2$ indicating the transition point. The Mott "lobe" is shown in Fig.1. Note that the kinetic energy and interaction energy are both zero in the Mott phase, which is an exact feature of the model.

Outside the Mott phase, the system is in a metallic phase described by four (two per spin species) dispersionless extremely correlated Fermi liquids (ECFLs) associated with the lower and upper Hubbard bands. Depending on the filling factor and value of U, for each spin species one can obtain a situation where only one kind of Fermi liquid appears to which there pertains a single well defined Fermi surface (see Fig. 1). For $n < 1$, this case physically resembles the ground state properties of the infinite U limit of the Hubbard model. The single Fermi liquid comes about because for small $n_\uparrow, n_\downarrow$ the lower Hubbard band is wide while the upper Hubbard band is very narrow. Even for $U < W/2$, although the two bands overlap, the upper band centered around U is too narrow to be occupied in the ground state. On the other hand, closer to half filling $n \leq 1$ for $U < W/2$ when the upper band can also be occupied, two coexistant Fermi liquids emerge, per spin species, associated with two Fermi surfaces. The boundary between these two cases is a Fermi surface topology changing transition of the Lifshitz type.

We now focus on the interaction driven Mott transition from the metallic phase. Unlike in the Gutwziller or slave boson methods, the hopping in the approach discussed here cannot be used as an indicator of the transition, as it remains constant. The density of doubly occupied sites $D = \langle \hat{D} \rangle / N$ (occupation of the upper Hubbard band) is a good parameter at half filling capturing the transition for this model. It takes a particularly simple form

$$D = \langle \hat{D} \rangle / N = \frac{1}{2} \int_{-W/2}^{-U} \rho(\epsilon)\, \mathrm{d}\epsilon \tag{6}$$

and depends only on U and the density of states. We show results for a few different density of states (DOSes)in Fig.2.

Note that an important feature of our Hubbard-I calculation is that the critical point does not depend on the lattice type. Furthermore, lattice dependent properties emerge for the number of doubly occupied sites. A linear dependence is associated here only with a rectangular density of states. In fact, the critical behaviour of D has universal properties depending only on spatial dimensions of the lattice. It is governed by the behaviour of the DOS at the bottom of the band. Indeed for parabolic energy profiles near the bottom of the band, as $\epsilon \to -W/2 \Rightarrow \Delta\epsilon_{\boldsymbol{k}} \propto \boldsymbol{k}^2$, we have for 1 dimension $\rho(-W/2 + \Delta\epsilon) \propto 1/\sqrt{\Delta\epsilon}$, 2 dimensions that $\rho(-W/2 + \Delta\epsilon) \propto$ const., and 3 dimensions $\rho(-W/2 + \Delta\epsilon) \propto \sqrt{\Delta\epsilon}$. Then we obtain that in 1d: $D \propto [(U_c - U)/U_c]^{\frac{1}{2}}$, 2d: $D \propto (U_c - U)/U_c$ while in 3d: $D \propto [(U_c - U)/U_c]^{\frac{3}{2}}$.



Figure 1: Ground state phase diagram in the $U - n$ plane. In all calculations we assume only nearest neighbour hopping $t_{ij} = -t$. A metallic ECFL exists for all $n \leq 1$. At low filling, the metal resembles the $U = \infty$ with no double occupancies. Closer to $n = 1$, the metal has double occupancies and is described by two Fermi liquids per spin species. In the H-I calculation, the boundary between these two behaviours is a Lifshitz-type transition, shown for hypercubic lattices in 1,2,3 dimensions and for the rectangular DOS. Shown is also the exact result for 1d, from [17] with rescaled $U \to U/2$, for comparison. All transition lines cross the $n = 1$ line at finite $U = W/2$ marking the point of Mott transition. Solid line depicts the Mott phase. The $n > 1$ can be obtained by invoking electron-hole symmetry. Inset: Ground state phase diagram in the $U - \mu$ plane. The electron hole symmetry line (dotted) separates $n < 1$ and $n > 1$ cases. The $n = 1$ case coincides with this line for $U < W/2$ whereas for $U > W/2$ opens into a Mott insulator "lobe" of finite width.

Relation to the Bond-charge Hubbard model While the results presented in the previous paragraph provide a clear and simple quantitative picture of a paramagnetic transition as well as the metallic phase, it is not a priori clear how good an approximation the Hubbard-I approach yields for the constrained model H_c. This model is related to the class of generalized Hubbard models, differing from the Hubbard Hamiltonian by an additional bond charge interaction term (see *e.g.* [13–15]):

$$H_{\mathrm{bond-charge}} = \hat{T} + U\hat{D} + \sum_{i,j,\sigma} X_{ij}(n_{i\bar{\sigma}} + n_{j\bar{\sigma}}) c^\dagger_{i\sigma} c_{j\sigma} \tag{7}$$

At a symmetry point, when the bond charge interaction is equal to the hopping $X_{ij} = -t_{ij}$, this generalized Hubbard model conserves the number of double occupancies. In fact, the resultant symmetric model can be mapped



Figure 2: The double occupancy per site D at half-filling $n = 1$ as a function of U. Shown are results for hypercubic lattices in 1, 2, 3 dimensions and for rectangular DOS. For all DOSes $D = 1/4$, at $U = 0$, as in the uncorrelated metal, although it is a ECFL here. The Mott transition takes place at $U = W/2$. Note different critical behavior in 1,2 and 3 d. The curve calculated for $1d$ coincides with known exact result [17] for rescaled $U \to U/2$.

onto the model H_c on bipartite lattices, via the canonical transform $U = \exp[-i\pi \sum_j \boldsymbol{R}_j n_{j\uparrow} n_{j\downarrow}]$, where $R_j = \pm 1$ on different sublattices, as shown in [16]. Analytical ground states of the symmetric bond-charge model were obtained in [15] in the regime $U > 2W$ at half filling, for arbitrary dimensions d. These results were improved upon in [17] revealing that a metal-insulator transition occurs at $U = W$, probably with ferromagnetic polarization on the metallic side (for $d > 1$), which is supported by numerical evidence [18].

The special case of $d = 1$, was exactly solved in [17] leading to the following main results: the metal to insulator transition occurs at $U_c = W$, and also the number of double occupancies is $D = 1/2\pi \mathrm{ArcCos}(U/U_c)$. In $d = 1$, the H-I approach used here underestimates the transition point, but remarkably the average number of double occupancies calculated for the 1-d DOS $\rho(\epsilon) = 1/(\pi\sqrt{1-\epsilon^2})$, using Eq.(6) leads to exactly the same function and prefactor as the exact solution (with $U_c = W/2$ now being the H-I critical value). This suggests, that the H-I works particularly well in 1 dimension. Furthermore, exact results away from half-filling $n = 1$ reveal, that there is a critical boundary separating states with doubly occupied sites from states with no such sites, which in the H-I framework considered here corresponds to the Lishitz lines U_L depicted in Fig.1. The exact boundary is given by the relation $U_L = W\cos(\pi(1-n))$ [17]. Notice that, apart from the value of the Mott transition point, the Lifshitz line obtained using the H-I approach here is in good quantitative agreement with these results (see Fig.1). In general dimensions, the expected phase diagram is expected to be qualitatively similar [19] as in 1 dimension. We thus see, that the phase diagram derived by the Hubbard-I approach is in good qualitative agreement with exact results.

Comparison with the Gutzwiller approach It is worthwhile to compare our H-I results, for half filling, with those obtained using the Gutzwiller approximation to the model H_c. We perform a standard calculation (see *e.g.* [12]), *i.e* assuming the reference state, that is Gutzwiller projected, to be the product of Fermi seas of the two spin species. Within the GA, the average energy of the Hamiltonian H_c can be written as:

$$\langle H_c \rangle = \sum_\sigma q_\sigma \epsilon_{0\sigma} + UD, \tag{8}$$

where $\epsilon_{0\sigma}$ is the energy of the Fermi sea of a given spin species. The band narrowing factor q_σ is easily found to be given by:

$$q_\sigma = \frac{(n_\sigma - D)(1 + D - n_\sigma - n_{\bar\sigma}) + D(n_{\bar\sigma} - D)}{n_\sigma(1 - n_\sigma)}. \tag{9}$$

Minimizing the average energy with respect to D, at half filling, we obtain a Gutzwiller-Brinkman-Rice transition(GBR) at a finite $U_c = -4|\epsilon_0|$, where ϵ_0 is the summed ground state energy of the noninteracting Fermi liquids. The GA density of doubles on the metallic side is linearly dependent on the interaction U, reducing to zero at the transition point:

$$D = \frac{1}{4}\Big(1 - U/U_c\Big), \quad U \leq U_c. \tag{10}$$

The GBR transition is of a quantitatively different character than the obtained H-I transition. Indeed, the distinct points are that the transition point is lattice dependent, while the metallic side has lattice independent double occupancies (which seem to be doubtful in the light of some exact results summarized above). As an aside, note however that, the choice of the reference state as noninteracting uncorrelated Fermi seas is of doubtful applicability, in the Gutzwiller analysis, of the model H_c given that it certainly does not even correspond to the ground state of a correlated hopping Hamiltonian anywhere, including in the limit $U = 0$. This is analogous to the Gutzwiller study of the bond-charge Hamiltonian Eq.(7) performed in [20], where the results for large bond charge interaction X cannot be considered reliable. Indeed, in particular at the symmetry point (which is of direct relevance to the model H_c), $X_{ij} = -t_{ij}$, Kollar and Vollhardt in [20] obtain a critical point $U_c = 0$ using the GA. Thus, rather interestingly, we observe that the Gutzwiller approximation yields a qualitatively better picture of the phase transition itself in the model H_c than in the bond-charge Hamiltonian. Finally, the GA shows a ferromagnetic instability in the bond-charge model, for certain lattices [20] before the Mott insulator transition. However, this is not the case for the GA in the H_c Hamiltonian. Indeed, as usual [6, 20] one can calculate the bulk magnetic susceptibility χ, which is here given by:

$$\chi^{-1} = \frac{1-(U/U_c)^2}{2\rho(\varepsilon_F)}(1-\rho(\varepsilon_F)U_c) \tag{11}$$

where $\rho(\varepsilon_F)$ is the density of states at the Fermi surface. Only one factor depends on U and can diverge here, accompanying the metal insulator transition $U \rightarrow U_c$. Thus, the GA, like the H-I calculation, does not describe a ferromagnetic metal before the Mott transition. However, apart from this the phases and critical points are better described by the H-I approach.

The Roth–corrected approach We have shown that the Hubbard-I approach to the model H_c gives the correct scale of energy for the Mott transition. For general dimensions, however, the critical point U_c does not depend on magnetic polarization, since the gap is always given by $U - W/2$. Recall, that in obtaining the description of the Mott transition we neglected terms, in the equations of motion, related to magnetic order and other terms. All these terms can be treated in a unified manner by the procedure invoked by Roth [21], which removes ambiguities in the decoupling of Green's functions equations of motion method. Applying the Roth procedure at the Hubbard-I level of equations of motion for the considered Hamiltonian H_c, we obtain the following Green's functions:

$$\Gamma^{\sigma}_{\boldsymbol{k}}(\omega) = \frac{1}{2\pi}\frac{n_{\bar{\sigma}}}{\omega - E_{\sigma}},\ \widetilde{\Gamma}^{\sigma}_{\boldsymbol{k}}(\omega) = \frac{1}{2\pi}\frac{h_{\bar{\sigma}}}{\omega - \widetilde{E}_{\sigma}} \tag{12}$$

where, the quasi-particle energy factors are

$$\begin{aligned} E_{\sigma} &= (U-\mu) + \epsilon_{\boldsymbol{k}}\langle n_{\boldsymbol{0}\bar{\sigma}} n_{\boldsymbol{\delta}\bar{\sigma}}\rangle / n_{\bar{\sigma}} \\ &-\epsilon_{\boldsymbol{k}}(\langle T_D\rangle + \langle T_X\rangle)/n_{\bar{\sigma}} + (\langle T_{UHB}\rangle - \langle T_{LHB}\rangle)/n_{\bar{\sigma}}, \quad (13) \\ \widetilde{E}_{\sigma} &= -\mu + \epsilon_{\boldsymbol{k}}\langle h_{\boldsymbol{0}\bar{\sigma}} h_{\boldsymbol{\delta}\bar{\sigma}}\rangle / h_{\bar{\sigma}} \\ &-\epsilon_{\boldsymbol{k}}(\langle T_D\rangle + \langle T_X\rangle)/h_{\bar{\sigma}} + (\langle T_{UHB}\rangle - \langle T_{LHB}\rangle)/h_{\bar{\sigma}}. \quad (14) \end{aligned}$$

The term $\langle T_X\rangle \equiv \langle c^{\dagger}_{\boldsymbol{0}\bar{\sigma}} c_{\boldsymbol{\delta}\bar{\sigma}} c^{\dagger}_{\boldsymbol{\delta}\sigma} c_{\boldsymbol{0}\sigma}\rangle$ describes the process of interchange of spins between neighbouring sites, while $\langle T_D\rangle \equiv \langle c^{\dagger}_{\boldsymbol{0}\bar{\sigma}} c_{\boldsymbol{\delta}\bar{\sigma}} c^{\dagger}_{\boldsymbol{0}\sigma} c_{\boldsymbol{\delta}\sigma}\rangle$ describes the process of doublon transfer. The averages in Eqs. (13, 14) are yet to be determined quantities, which is a standard feature of Roth's method. Notice, that disregarding the second lines in Eqs. (13, 14), and considering site occupations as uncorrelated, one recovers the Hubbard I results.

A full analysis of the Roth solutions shall be considered elsewhere [22], while here we only consider implications for the Mott phase in which $\langle T_X\rangle, \langle T_D\rangle, \langle T_{UHB}\rangle, \langle T_{LHB}\rangle$ vanish identically for the considered model H_c. The closure of the spectral gap to excitations indicates the Mott phase instability. Consider the case half-filled case with $n_{\uparrow} = n_{\downarrow} = 1/2$. If there are no intersite correlations in the Mott states, $U_c = W/2$ as indicated above. On the other hand for saturated ferromagnetic correlation, *i.e.* macroscopic separation of the system into two ferromagnetic domains, Eqs. (13, 14) show that there is no renormalization of the band width in the Hubbard bands, and thus $U_c = W$. As all Mott states have the same energy, this analysis indicates that the Mott phase is unstable already at $U_c = W$ (which is in good agreement with known results summarized earlier), and changes probably into a ferromagnetic metallic state.

Conclusions and Outlook In this work, we have analyzed the Hubbard I approximation, explicitly showing that its known drawbacks originate from the interband hopping $T_{\pm 1}$ terms in the Hubbard model. We proposed to use the H-I approach in conjunction with perturbational expansion. The H-I approach, in combination with lowest order perturbational expansion, leads to a physically appealing, picture of the Mott transition including the appearance of a extremely correlated Fermi liquid in its vicinity, which is complementary to the Gutzwiller-Brinkman-Rice picture.

It is natural to wonder how well this approach fits as a description of Mott transitions and the surrounding ECFL in realistic strongly correlated electron models. In this regard, we emphasize that the double occupancy conserving Hamiltonian H_c analyzed here is equivalent to the Hubbard model with bond charge interactions at the symmetry point $X = -t$ – which has been argued to be a quite realistic value (see *e.g.* [15, 17]). Our H-I or Roth improved H-I calculations compare quite favourably with known and expected results for the latter model, and are remarkably consistent with them in 1 dimension. Indeed, one would expect the picture presented here to hold in the neighbourhood of the symmetry point. On the other hand, for pure Hubbard like–systems, *i.e.* X close to 0, of course this strong coupling picture of the Mott transition can only be qualitatively correct (when antiferromagnetic order is suppressed) and that too only near the transition point. This gives rise to an open question concerning the possibility of obtaining an interpolating scheme between these two extreme cases.

In this regard, we propose to use the following variational ground state for systems exhibiting metal to Mott insulator transitions:

$$|\Psi\rangle = \exp[-\alpha\hat{T}_0]\exp[-\eta\hat{D}]|\Psi_0\rangle = \exp[-\alpha\hat{T}_0]|\mathrm{GWF}\rangle, \tag{15}$$

where α, η are variational parameters and $|\Psi_0\rangle$ is an appropriate reference state. This state is an extension of the standard Gutzwiller Wave Function (GWF) and should provide an improved description of the metallic side of the transition. Note that the two exponential terms commute and together form the exponential of $-\alpha H_c(U = \eta/\alpha)$, which may be viewed as a partial projection on to the ground state of $H_c(U = \eta/\alpha)$. This makes a connection with the ECFL properties described in this paper and allows for dressing of the GWF with precursors of the Hubbard bands. It is worth mentioning here that a similar function (containing hopping only of the lower Hubbard band in T_0) has already been used in [23] and has provided excellent results in comparison with exact results for 2 electrons on the Hubbard square.

Acknowledgements We thank Roman Micnas for many interesting discussions.

[1] N. F. Mott, Proc. Phys. Soc. London, **A 63** 416 (1949); N. F. Mott, Philos. Mag. **6**, 287 (1961).

[2] J. Hubbard, Proc. Roy. Soc. London **276**, 238 (1963).

[3] J. Hubbard, Proc. Roy. Soc. London **281**, 401 (1964).

[4] D. M. Edwards and A. C. Hewson, Rev. Mod. Phys. **40**, 810 (1968).

[5] M. Gutzwiller, Phys. Rev. 137, A1726 (1965).

[6] W. F. Brinkman and T. M. Rice, Phys. Rev. **B 2**, 4302 (1970).

[7] A. Georges, G. Kotliar, W. Krauth, M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996) .

[8] D. Hansen, E. Perepelitsky, B. S. Shastry, Phys. Rev. B **83**, 205134 (2011).

[9] A. B. Harris and R. V. Lange, Phys. Rev. 157, 295 (1967).

[10] K. A. Chao, J. Spałek and A. M. Oleś, Phys. Rev. B 18, 3453 (1978); A. H. MacDonald, S. M. Girvin and Yoshioka, Phys. Rev. B 37, 9753 (1988).

[11] B. S. Shastry, Phys. Rev. B **81** 045121 (2010).

[12] Dieter Vollhardt, Rev. Mod. Phys. **56**, 99 (1984).

[13] R. Micnas, J. Ranninger, S. Robaszkiewicz, Phys. Rev. B **39**, 11653 (1989).

[14] F. Marsiglio and J. E. Hirch Phys. Rev. B **41**, 6435 (1990).

[15] R. Strack and D. Vollhardt, Phys. Rev. Lett. 70, 2637 (1993).

[16] F. Gebhard, K. Bott, M. Scheidler, P. Thomas, S. W. Koch, Phil. Mag. **B** 75, pp. 13-46 (1997).

[17] A. A. Ovchinnikov, Mod. Phys. Lett. B **7**, 1397 (1993); A. A. Ovchinnikov, J. Phys.: Condens. Matter 6, (1994).

[18] E. R. Gagliano, A. A. Aligia, L. Arrachea, M Avignon, Phys. Rev. **51** 14012 (1994).

[19] A. Schadschneider, Phys. Rev. B **51**, 10386 (1995).

[20] M. Kollar, D. Vollhardt, Phys. Rev. B **63**, 045107 (2001).

[21] L. M. Roth, Phys. Rev. Lett. **25**, 1431 (1968); L. M. Roth, Phys. Rev. **184**, 451(1969).

[22] P. R. Grzybowski, R. W. Chhajlany, *in preparation*

[23] D. Baeriswyl, D. Eichenberger, M. Menteshashvili, New. J. Phys. **11**, 075010 (2009).